

ARITHMETIC-GEOMETRIC MEANS FOR HYPERELLIPTIC CURVES AND CALABI-YAU VARIETIES

KEIJI MATSUMOTO AND TOMOHIDE TERASOMA

Dedicated to Professor Kazuo Okamoto on the occasion of his 60-th birthday.

ABSTRACT. In this paper, we define a generalized arithmetic-geometric mean μ_g among 2^g terms motivated by 2τ -formulas of theta constants. By using Thomae's formula, we give two expressions of μ_g when initial terms satisfy some conditions. One is given in terms of period integrals of a hyperelliptic curve C of genus g . The other is by a period integral of a certain Calabi-Yau g -fold given as a double cover of the g -dimensional projective space $\mathbf{P}^g$.

1. INTRODUCTION

Let $\{a_{n,0}\}_n$ and $\{a_{n,1}\}_n$ be positive real sequences defined by the recurrence relations

$$(1.1) \quad a_{n+1,0} = \frac{a_{n,0} + a_{n,1}}{2}, \quad a_{n+1,1} = \sqrt{a_{n,0}a_{n,1}},$$

and initial terms $a_{0,0} = a_0$, $a_{0,1} = a_1$ with $0 < a_1 < a_0$. One can easily show that $\{a_{n,0}\}_n$ and $\{a_{n,1}\}_n$ have a common limit, which is called the arithmetic-geometric mean of a_0 and a_1 , and is denoted by $\mu_1(a_0, a_1)$. By the homogeneity of the arithmetic and geometric means, we have $\mu_1(ca_0, ca_1) = c\mu_1(a_0, a_1)$ for any positive real number c .

On the other hand, two Jacobi's theta constants θ_0 and θ_1 satisfy the following 2τ -formulas:

$$(1.2) \quad \theta_0(2\tau)^2 = \frac{\theta_0(\tau)^2 + \theta_1(\tau)^2}{2}, \quad \theta_1(2\tau)^2 = \theta_0(\tau)\theta_1(\tau),$$

where

$$\theta_i(\tau) = \sum_{n \in \mathbf{Z}} \exp(\pi \sqrt{-1}(n^2\tau + in)), \quad i = 0, 1,$$

and τ belongs to the upper half space $\mathbf{H}$. If we find an element $\tau \in \mathbf{H}$ such that $\theta_1(\tau)^2/\theta_0(\tau)^2 = a_1/a_0$ for given initial terms a_0 and a_1 , then we have

$$\frac{a_0}{\mu_1(a_0, a_1)} = \frac{\theta_0(\tau)^2}{\mu_1(\theta_0(\tau)^2, \theta_1(\tau)^2)} = \frac{\theta_0(\tau)^2}{\mu_1(\theta_0(2^n\tau)^2, \theta_1(2^n\tau)^2)} = \theta_0(\tau)^2$$

2000 *Mathematics Subject Classification.* Primary 14K20; Secondary 32G20.

by (1.1), (1.2) and $\lim_{n \rightarrow \infty} \theta_i(2^n \tau) = 1$. Moreover, Jacobi's formula between $\theta_0(\tau)^2$ and an elliptic integral implies that

$$\frac{a_0}{\mu_1(a_0, a_1)} = \frac{2}{\pi} \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}, \quad k = \frac{\sqrt{a_0^2 - a_1^2}}{a_0}.$$

In this paper, we define a generalized arithmetic-geometric mean μ_g among 2^g terms $(\dots, a_I, \dots)$ ($I \in \mathbf{F}_2^g$) motivated by the 2τ -formulas (2.3) of theta constants obtained by Theorem 2 in [3] p.139. By using Thomae's formula, we give two expressions of μ_g whose initial terms are given as (3.1) for some $2g+1$ real numbers p_j . One is given in terms of period integrals of the hyperelliptic curve C of genus g represented by the double cover of the complex projective line $\mathbf{P}^1$ branching at ∞ and $2g+1$ points p_j . The other is by a period integral of the Calabi-Yau g -fold which is the double cover of the g -dimensional projective space $\mathbf{P}^g$ branching along the dual hyperplanes of the images of ∞ and p_j ($j = 1, \dots, 2g+1$) under the Veronese embedding of $\mathbf{P}^1$ into $\mathbf{P}^g$.

In 1876, Borchardt studied in [1] the case of $g = 2$: the generalized arithmetic-geometric mean μ_2 of $a = (a_{00}, a_{01}, a_{10}, a_{11})$ was given by the iteration of four means

$$\begin{aligned} & \frac{a_{00} + a_{01} + a_{10} + a_{11}}{4}, \quad \frac{\sqrt{a_{00}a_{01}} + \sqrt{a_{11}a_{10}}}{2}, \\ & \frac{\sqrt{a_{00}a_{10}} + \sqrt{a_{11}a_{01}}}{2}, \quad \frac{\sqrt{a_{00}a_{11}} + \sqrt{a_{10}a_{01}}}{2}, \end{aligned}$$

and $\mu_2(a)$ was expressed in terms of period integrals of a hyperelliptic curve of genus 2. Mestre showed in [4] that $\mu_2(a)$ could be expressed in terms of μ_1 and some algebraic functions of a when

$$a_{00} > a_{01} = a_{10} > a_{11}, \quad a_{00}a_{11} > a_{01}a_{10}.$$

2. COMPARISON TO THETA CONSTANTS

We define a hyperelliptic curve C of genus g by

$$C : y^2 = (x - p_1) \cdots (x - p_{2g+1}),$$

where p_j 's are real numbers satisfying $p_1 < \cdots < p_{2g+1}$. As in [6] p.76, we choose the cycles $A_1, \dots, A_g, B_1, \dots, B_g$ in the union of the following two sheets (I), (II) in Figure 1. Here $\mathbf{R}_+$ is the set of non-negative real numbers, the range of values of y is written, and the cycles in the sheet II are written in thick lines. Note that the cycles satisfy

$$A_i \cdot A_j = B_i \cdot B_j = 0, \quad A_i \cdot B_j = \delta_{ij}$$

for $1 \leq i, j \leq g$ under the intersection form.

We define holomorphic forms ω_j for $j = 1, \dots, g$ as

$$\omega_j = \frac{x^{j-1} dx}{y}.$$

Sheet I (solid line)

$$p_1 \xrightarrow[\mathbf{R}_+]{-\mathbf{R}_+} p_2 \xrightarrow[\sqrt{-1} \cdot \mathbf{R}_+]{\mathbf{R}_+} p_3 \xrightarrow[-\mathbf{R}_+]{\mathbf{R}_+} p_4 \xrightarrow[-\sqrt{-1} \cdot \mathbf{R}_+]{\mathbf{R}_+} \dots p_{2g+1} \xrightarrow{\mathbf{R}_+} \infty$$

Sheet II (dotted line)

$$p_1 \xrightarrow[-\mathbf{R}_+]{\mathbf{R}_+} p_2 \xrightarrow[-\sqrt{-1} \cdot \mathbf{R}_+]{-\mathbf{R}_+} p_3 \xrightarrow[\mathbf{R}_+]{-\mathbf{R}_+} p_4 \xrightarrow[\sqrt{-1} \cdot \mathbf{R}_+]{-\mathbf{R}_+} \dots p_{2g+1} \xrightarrow{\mathbf{R}_+} \infty$$

Cycles




FIGURE 1. Symplectic basis

We define integrals $T_i^{(j)}$ by

$$T_i^{(j)} = \int_{p_i}^{p_{i+1}} \frac{x^{j-1} dx}{\sqrt{\prod_{k=1}^i (x - p_k) \prod_{k=i+1}^{2g+1} (p_k - x)}}$$

for $1 \leq i \leq 2g$ and $1 \leq j \leq g$. Then the integrals $T_i^{(j)}$ are positive real numbers. Using these integrals, we express the period integrals of C :

$$\int_{A_i} \omega_j = (-1)^i 2T_{2i-1}^{(j)}, \quad \int_{B_i} \omega_j = 2\sqrt{-1} \left(\sum_{k=i}^g (-1)^{k+1} T_{2k}^{(j)} \right).$$

We set

$$(2.1) \quad A = \left(\int_{A_i} \omega_j \right)_{ij}, \quad B = \left(\int_{B_i} \omega_j \right)_{ij}$$

and consider the normalized period matrix τ by A-period:

$$(2.2) \quad \tau = BA^{-1}.$$

By Riemann's bilinear relations, $\det(A)$ is a non-zero real number and τ is a symmetric matrix whose imaginary part is positive definite. Note also that τ is purely imaginary.

Remark 2.1. Since the Vandermonde matrix $\det(x_i^{j-1})_{1 \leq i, j \leq g}$ is positive on $p_{2i-1} \leq x_i \leq p_{2i}$, $(-1)^{g(g+1)/2} \det(A)$ is positive.

For $I = (i_1, \dots, i_g) \in \mathbf{F}_2^g$, we define theta constants as

$$\theta_I(\tau) = \sum_{n \in \mathbf{Z}^g} \exp(\pi\sqrt{-1} \cdot n \tau \cdot {}^t n + \pi\sqrt{-1} \cdot n \cdot {}^t I).$$

Proposition 2.2. *Let M be a positive definite symmetric $g \times g$ real matrix. Then $\theta_I(\sqrt{-1}M)$ is positive for each $I \in \mathbf{F}_2^g$.*

Proof. By the inversion formula of the theta function in [5] p.195, we have

$$\sqrt{\det(M)} \cdot \theta_I(\sqrt{-1}M) = \sum_{n \in \mathbf{Z}^g} \exp\left(\sqrt{-1}\pi\left(n + \frac{I}{2}\right)(\sqrt{-1}M^{-1}) \cdot \left(n + \frac{I}{2}\right)\right),$$

where $\sqrt{\det(M)}$ takes a positive value. Since each term of the right hand side is positive, the left hand side is positive. $\square$

We consider variable $u = (u_I)_{I \in \mathbf{F}_2^g}$ whose coordinates are indexed by $\mathbf{F}_2^g$. The pair $(\theta_I(\tau))_I$ is denoted by $\theta(\tau)$. For $I \in \mathbf{F}_2^g$, we define quadratic polynomials $F_I(u)$ of 2^g variables $u = (u_I)_{I \in \mathbf{F}_2^g}$ by

$$F_I(u) = \frac{1}{2^g} \sum_{P \in \mathbf{F}_2^g} u_{I+P} u_P.$$

We remark that the coefficients of $2^g F_I(u)$ are in $\mathbf{Z}_{\geq 0}$. By Theorem 2 in [3] p.139, we have 2τ -formulas of theta constants

$$(2.3) \quad \theta_I(2\tau)^2 = F_I(\theta(\tau))$$

for $I \in \mathbf{F}_2^g$.

Now prepare some combinatorial notations for the statement of Thomae's formula. For an index $I \in \mathbf{F}_2^g$, we define a subset S_I of $R = \{1, \dots, 2g+1, \infty\}$ as follows. Let η_i be elements of $M(2, g, \mathbf{F}_2)$ defined as

$$\eta_{2i-1} = \begin{pmatrix} 0 & \cdots & 0 & \overset{i\text{-th}}{1} & 0 & \cdots & 0 \\ 1 & \cdots & 1 & 0 & 0 & \cdots & 0 \end{pmatrix},$$

$$\eta_{2i} = \begin{pmatrix} 0 & \cdots & 0 & \overset{i\text{-th}}{1} & 0 & \cdots & 0 \\ 1 & \cdots & 1 & 1 & 0 & \cdots & 0 \end{pmatrix},$$

for $i = 1, \dots, 2g+1$. Then a subset T_I of $R - \{2g+1, \infty\} = \{1, 2, \dots, 2g\}$ is characterized by the equality

$$\binom{0}{I} = \sum_{j \in T_I} \eta_j.$$

We set

$$S_I = \begin{cases} T_I & \text{if } \#T_I \text{ is even,} \\ T_I \cup \{2g+1\} & \text{if } \#T_I \text{ is odd.} \end{cases}$$

Let U be the set $\{1, 3, 5, \dots, 2g+1\}$ and $R_1 \circ R_2$ be the symmetric difference of sets R_1 and R_2 .

Proposition 2.3 ([6] p.120, [2]). *Let A be the period matrix of C in (2.1). Then we have*

$$(2.4) \quad \frac{(2\pi)^{2g}\theta_I(\tau)^4}{\det(A)^2} = \prod_{i < j, i, j \in S_I \circ U} (p_j - p_i) \prod_{i < j, i, j \notin S_I \circ U} (p_j - p_i).$$

Here we used the fact that $\theta_I(\tau)$ is a real number to determine the sign of Thomae's formula in [6].

3. STATEMENT AND PROOF OF THE MAIN THEOREM

Definition 3.1 (AGM sequences).

- (1) For an element $u = (u_I)_I \in \mathbf{R}_+^{2g}$, we define the termwise root $\sqrt{u}$ of u by $(\sqrt{u_I})_I$.
- (2) Let $a = (a_I)_I$ be an element in $(\mathbf{R}_+)^{2g}$. We define $a_k = (a_{k,I})_I$ inductively by the relation

$$a_{0,I} = a_I, \quad a_{k+1,I} = F_I(\sqrt{a_k}).$$

A proof of the following proposition will be left to readers.

Proposition-Definition 3.2 (Generalized arithmetic-geometric mean). *For an element $a = (a_I)_I$ in $(\mathbf{R}_+)^{2g}$, the limits $\lim_{k \rightarrow \infty} a_{k,I}$ exist and are independent of indexes I . This common limit is called the generalized arithmetic-geometric mean of $a = (a_I)_I$ and denoted by $\mu_g(a)$.*

Problem 3.3. *Is it possible to express the generalized arithmetic-geometric mean $\mu_g(a)$ of $a = (a_I)_I \in (\mathbf{R}_+)^{2g}$ in terms of period integrals of a family of varieties parametrized by a ?*

Theorem 3.4. *Let $p_1 < \dots < p_{2g+1}$ be real numbers. We define a_I by*

$$(3.1) \quad a_I = \sqrt{\prod_{i < j, i, j \in S_I \circ U} (p_j - p_i) \prod_{i < j, i, j \notin S_I \circ U} (p_j - p_i)}.$$

Then we have

$$\mu_g(a) = \frac{(2\pi)^g}{|\det(A)|},$$

where A is the period matrix of C in (2.1).

Proof. By the initial condition, we have

$$a_{0,I} = \frac{(2\pi)^g \theta_I(\tau)^2}{|\det(A)|}.$$

We show that

$$a_{n,I} = \frac{(2\pi)^g \theta_I(2^n \tau)^2}{|\det(A)|},$$

by induction on n . Since $\theta_I(2^n \tau)$ is a positive real number by Proposition 2.2 for each I , we have

$$\begin{aligned} a_{n+1,I} &= F(\sqrt{a_n}) \\ &= \frac{(2\pi)^g \cdot F(\theta(2^n \tau))}{|\det(A)|} && \text{(by the induction hypothesis)} \\ &= \frac{(2\pi)^g \cdot \theta_I(2^{n+1} \tau)^2}{|\det(A)|} && \text{(by the formula (2.3))} \end{aligned}$$

Therefore we have

$$\lim_{n \rightarrow \infty} a_{n,I} = \frac{(2\pi)^g}{|\det(A)|}.$$

□

4. PERIOD OF CALABI-YAU VARIETY OF CERTAIN TYPE

We study a relation between the generalized arithmetic-geometric mean of the last section and a period of a Gorenstein Calabi-Yau variety of a certain type.

Definition 4.1 (Calabi-Yau varieties). *A variety X only with Gorenstein singularities is called a Calabi-Yau variety if the dualizing sheaf of X is trivial and X has a global crepant resolution.*

Let $\mathbf{P} = \mathbf{P}^g$ be the g dimensional projective space and $H_1 \cdots H_{2g+2}$ be hyperplanes of $\mathbf{P}$. There is a unique line bundle $\mathcal{L}$ on $\mathbf{P}$ and a unique isomorphism $\varphi : \mathcal{L}^{\otimes 2} \simeq \mathcal{O}_X(-\sum_{i=1}^{2g+2} H_i)$ up to a non-zero constant. Using the isomorphism φ , we define a double covering $X = \text{Spec}(\mathcal{O}_X \oplus \mathcal{L})$, where the multiplication on $\mathcal{L} \otimes \mathcal{L} \rightarrow \mathcal{O}_X$ is given by the isomorphism φ .

By the following Proposition 4.2, X becomes a Calabi-Yau variety, since it admits a global crepant resolution.

Proposition 4.2.

- (1) *If $\cup_{i=1}^{2g+2} H_i$ is normal crossing, then the variety X has only Gorenstein singularities. Also it admits a global crepant resolution.*
- (2) *Under the above hypotheses, the dualizing sheaf is isomorphic to the structure sheaf.*

Proof. (1) We choose a local coordinate $\xi_1, \dots, \xi_g$ of $\mathbf{P}$ such that the divisor $\cup_{i=1}^{2g+2} H_i$ is locally defined by $\xi_1 \cdots \xi_h = 0$ ($h \leq g$). Then the variety X is locally defined by the equation $\eta^2 = \xi_1 \cdots \xi_h$. Therefore X is locally isomorphic to $\text{Spec}(\sigma \cap M^*) \times \mathbf{A}^{g-h}$, where

$$M^* = \mathbf{Z}^h + \left(\frac{1}{2}, \dots, \frac{1}{2}\right) \mathbf{Z} \subset \mathbf{Q}^h, \quad \sigma = (\mathbf{R}_+)^h.$$

Let σ be the dual simplex of σ and M be the dual lattice of M^* . Since σ is generated by elements contained primitive hyperplanes, X is Gorenstein. We can construct a global crepant resolution as follows. We make a refinement of the simplex σ into a regular fan $\cup_{\mathbf{w} \in \rho_h} \sigma_{\mathbf{w}}$ indexed by the set ρ_h of “unfair

tournament" of $\{1, \dots, h\}$. A sequence $\mathbf{w} = (w_1, \dots, w_{h-1})$ is an element of the set ρ_h if it satisfies the following properties:

- (i) w_1 is equal to 1 or 2 and
- (ii) w_i is equal to w_{i-1} or $i+1$ for $2 \leq i \leq h-1$.

For an element $\mathbf{w}$ of ρ_h , we define $\sigma_{\mathbf{w}}$ as a cone generated by

$$B_{\mathbf{w}} = \{u_1 = e_1 + e_2, u_2 = e_{w_1} + e_3, u_3 = e_{w_2} + e_4 \dots, u_{h-1} = e_{w_{h-2}} + e_h, \\ u_h = 2e_{w_{h-1}}\},$$

where e_i is the standard basis of $\mathbf{Z}^h \supset M$. Since the set $B_{\mathbf{w}}$ is a free base of M , the fan $\cup_{\mathbf{w} \in \rho_h} \sigma_{\mathbf{w}}$ is regular and it defines a smooth toric variety $\tilde{X}$. The coordinates associated to $\mathbf{Z}^h \subset M^*$ are written as $\xi_1, \dots, \xi_h$. (η corresponds to $\frac{1}{2}(1, \dots, 1)$.) Let $z_1, \dots, z_h$ be the coordinates associated to the dual base $B_{\mathbf{w}}$ of M . Then we have

$$z_1^{u_1} \dots z_h^{u_h} = \xi_1^{e_1} \dots \xi_h^{e_h}.$$

Thus $\xi_1^{\frac{1}{2}} \dots \xi_h^{\frac{1}{2}} = z_1 \dots z_h$. Let ω_X be the rational differential form on X defined by

$$\omega_X = \xi_1^{-\frac{1}{2}} \dots \xi_h^{-\frac{1}{2}} d\xi_1 \wedge \dots \wedge d\xi_h \wedge d\xi_{h+1} \wedge \dots \wedge d\xi_g.$$

It is a generator of the dualizing sheaf of X . The pull back of ω_X to the affine toric variety associated to $\sigma_{\mathbf{w}}$ is a non-zero constant multiple of

$$dz_1 \wedge \dots \wedge dz_h \wedge d\xi_{h+1} \wedge \dots \wedge d\xi_g.$$

Therefore the map $\tilde{X} \times \mathbf{A}^{g-h} \rightarrow X$ is a crepant resolution. Since the local crepant resolutions depend only on the choice of order of the components of the branching divisor, they are patched together into a global crepant resolution.

(2) In this proof, we use symbols $\xi_1, \dots, \xi_g$ as the global inhomogeneous coordinates of $\mathbf{P}$ for the infinite hyperplane H_{2g+2} . Let $l_i = l_i(\xi)$ be an inhomogeneous linear form defining the hyperplane H_i for $i = 1, \dots, 2g+1$. Then a defining equation of the double covering X can be written as

$$\eta^2 = \prod_{i=1}^{2g+1} l_i(\xi).$$

As is shown in the proof of (1),

$$(4.1) \quad \omega_X = \frac{1}{\eta} d\xi_1 \wedge \dots \wedge d\xi_g$$

is a global generator of the dualizing sheaf of X . □

For real numbers $p_1 < \dots < p_{2g+1}$, we define linear forms l_i by

$$l_i = \xi_1 - p_i \xi_2 + p_i^2 \xi_3 + \dots + (-1)^{g-1} p_i^{g-1} \xi_g + (-1)^g p_i^g$$

and set $H_i = \{l_i = 0\}$. By using the Vandermonde matrix, we see that $\cup_{i=1}^{2g+2} H_i$ is a normal crossing divisor.

We define a subset Δ of $\mathbf{R}^g$ as

$$\Delta = \{(x_1, \dots, x_g) \mid (-1)^{i-1} l_{2i-1}(x_1, \dots, x_g) \geq 0 \text{ for } i = 1, \dots, g+1, \text{ and} \\ (-1)^i l_{2i}(x_1, \dots, x_g) \geq 0 \text{ for } i = 1, \dots, g\}.$$

We set

$$\omega_X = \frac{1}{\eta} d\xi_1 \wedge \dots \wedge d\xi_g, \text{ and } \gamma_{\pm} = \{(\xi, \eta) \in X \mid \xi \in \Delta, \pm\eta \geq 0\}.$$

Then $\gamma = \gamma_+ - \gamma_-$ defines a g -chain in X . We have the following relation between the generalized arithmetic-geometric mean and a period of the Calabi-Yau variety X . The following theorem is obtained by Theorem 2 in [7].

Theorem 4.3. *Let $a = (a_I)_I$ be an element of $\mathbf{R}_+^g$ defined in (3.1). Under the above notation, we have*

$$\mu_g(a) = \frac{2\pi^g}{\int_{\gamma} \omega_X}.$$

Proof. Let C_j be a copy of the curve C given by $y_j^2 = \prod_{i=1}^{2g-1} (x_j - p_i)$. We define a map $\pi : C_1 \times \dots \times C_g \rightarrow X$ by sending $((x_1, y_1), \dots, (x_g, y_g))$ to the point whose ξ_k -coordinate and η -coordinate are the $(g+1-k)$ -th elementary symmetric function of $x_1, \dots, x_g$ and $\prod_{j=1}^g y_j$, respectively. Then we have

$$\pi^* \omega_X = \sum_{\sigma \in \mathcal{S}_g} \text{sgn}(\sigma) \boxtimes_{i=1}^g \omega_{\sigma(i)}.$$

Since $\pi_*(A_1 \times \dots \times A_g) = (-1)^{g(g+1)/2} 2^{g-1} \gamma$, we have

$$2^{g-1} \int_{\gamma} \omega_X = |\det(A)|.$$

By Theorem 3.4, we have the theorem. $\square$

5. GENUS TWO CASE

In this section, we will give a detailed study for the case of $g = 2$. Refer to [1] and [4] for the original results by Borchartd and recent related works by Mestre, respectively. We begin with $(a_{00}, a_{01}, a_{10}, a_{11})$ as initial data for AGM sequences. The recursive relations for $a_{k,I}$ ($I \in \mathbf{F}_2^2, k = 0, 1, \dots$) are given as $a_{0,I} = a_I$ and $a_{k+1,I} = F_I(\sqrt{a_{k,00}}, \dots, \sqrt{a_{k,11}})$, where

$$\begin{aligned} F_{00}(u_{00}, u_{01}, u_{10}, u_{11}) &= \frac{1}{4}(u_{00}^2 + u_{01}^2 + u_{10}^2 + u_{11}^2), \\ F_{01}(u_{00}, u_{01}, u_{10}, u_{11}) &= \frac{1}{2}(u_{00}u_{01} + u_{11}u_{10}), \\ F_{10}(u_{00}, u_{01}, u_{10}, u_{11}) &= \frac{1}{2}(u_{00}u_{10} + u_{11}u_{01}), \\ F_{11}(u_{00}, u_{01}, u_{10}, u_{11}) &= \frac{1}{2}(u_{00}u_{11} + u_{10}u_{01}). \end{aligned}$$

In the following, we assume that $a_{00} > a_{10} > a_{11} > a_{01}$ and $a_{00}a_{01} > a_{10}a_{11}$. First we define positive real numbers $k_1 > k_2$ and $0 < l_2 < l_1 < 1$ such that

$$(a_{00} + a_{01})^2 - (a_{10} + a_{11})^2 = k_1^2, \quad (a_{00} - a_{01})^2 - (a_{10} - a_{11})^2 = k_2^2,$$

$$\begin{aligned} a_{00} + a_{01} &= \frac{1 + l_1^2}{1 - l_1^2} k_1, & a_{10} + a_{11} &= \frac{2l_1}{1 - l_1^2} k_1, \\ a_{00} - a_{01} &= \frac{1 + l_2^2}{1 - l_2^2} k_2, & a_{10} - a_{11} &= \frac{2l_2}{1 - l_2^2} k_2, \end{aligned}$$

We set

$$\begin{aligned} p_1 &= 0, & p_2 &= \frac{1}{(1 - l_2^2)(1 - l_1^2)}, \\ p_3 &= \frac{2(l_1 l_2 + 1) a_{00}}{(1 - l_1^2)(1 - l_2^2)(k_1 + k_2)(1 - l_1 l_2)}, \\ p_4 &= \frac{2(l_1 l_2 + 1) a_{01}}{(1 - l_1^2)(1 - l_2^2)(k_1 - k_2)(1 - l_1 l_2)}, \\ p_5 &= \frac{4a_{00}a_{01}}{(k_1 - k_2)(k_1 + k_2)(1 - l_2^2)(1 - l_1^2)}. \end{aligned}$$

Then we have

$$\begin{aligned} (5.1) \quad (a_{00}^2 : a_{01}^2 : a_{10}^2 : a_{11}^2) &= ((p_3 - p_1)(p_5 - p_1)(p_5 - p_3)(p_4 - p_2) : \\ &\quad (p_4 - p_1)(p_5 - p_1)(p_5 - p_4)(p_3 - p_2) : \\ &\quad (p_3 - p_2)(p_5 - p_2)(p_5 - p_3)(p_4 - p_1) : \\ &\quad (p_4 - p_2)(p_5 - p_2)(p_5 - p_4)(p_3 - p_1)). \end{aligned}$$

Therefore by Theorem 3.4, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} a_{n,00} &= \frac{4\pi^2 a_{00}}{|\det(A)| \sqrt{(p_3 - p_1)(p_5 - p_1)(p_5 - p_3)(p_4 - p_2)}} \\ &= \frac{8\pi^2}{|\det(A)|} \cdot (1 - l_1^2)^2 (1 - l_2^2)^2 \sqrt{\frac{(a_{00}a_{01} - a_{10}a_{11})^3 (1 - l_1 l_2)^3}{a_{00}a_{01}a_{10}a_{11}(l_1^2 - l_2^2)(1 + l_1 l_2)}}. \end{aligned}$$

where A is the period matrix of C in (2.1).

Using the result of §4, we have

$$|\det(A)| = 4 \cdot \int_{\Delta} \frac{d\xi_1 \wedge d\xi_2}{\sqrt{\prod_{i=1}^5 (\xi_1 - p_i \xi_2 + p_i^2)}},$$

where Δ is a domain in $\mathbf{R}^2$ defined by $l_1 \geq 0, -l_2 \geq 0, -l_3 \geq 0, l_4 \geq 0$ and $l_5 \geq 0$. This is a period integral of the covering X of $\mathbf{P}^2$ defined by

$$\eta^2 = \prod_{i=1}^5 (\xi_1 - p_i \xi_2 + p_i^2).$$

We notice that the variety X is the (nodal) Kummer surface of the Jacobian of C .

Remark 5.1. *When*

$$a_{00} > a_{01} = a_{10} > a_{11}, \quad a_{00}a_{11} > a_{01}a_{10},$$

$\mu_2(a)$ can be expressed in terms of the arithmetic-geometric mean μ_1 and expressions $p_2, \dots, p_5$ by a (see [4]).

REFERENCES

- [1] Borchardt, C.W.: Über das arithmetisch-geometrische Mittel aus vier Elementen, *Berl. Monatsber.*, 53 (1876), 611–621.
- [2] Fay, J.: *Theta functions on Riemann surfaces*, Lecture note in Math 352. Springer, Berlin-New York, 1973.
- [3] Igusa, J.: *Theta functions*, Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen 194, Springer-Berlin-Heidelberg, New York, 1972.
- [4] Mestre, J.: Moyenne de Borchardt et integrales elliptiques, *C. R. Acad. Sci. Paris Ser. I Math.* 313 (1991), no. 5, 273–276.
- [5] Mumford, D.: *Tata lectures on Theta I*, progress in Math 28. Birkhäuser, Boston-Basel-Berlin, 1983.
- [6] Mumford, D.: *Tata lectures on Theta II*, progress in Math 43. Birkhäuser, Boston-Basel-Berlin, 1984.
- [7] Terasoma, T.: Exponential Kummer coverings and determinants of hypergeometric functions. *Tokyo J. Math.* 16 (1993), no. 2, 497–508.

Keiji MATSUMOTO

Department of Mathematics
Hokkaido University
Sapporo, 060-0810, Japan
e-mail: matsu@math.sci.hokudai.ac.jp

Tomohide TERASOMA

Graduate School of Mathematical Sciences
The University of Tokyo
Komaba, Meguro, Tokyo, 153-8914, Japan
e-mail: terasoma@ms.u-tokyo.ac.jp